

References

1. McCrindle B, Rowley A, Newburger J, et al. Diagnosis, treatment, and long-term management of kawasaki disease: A scientific statement for health professionals from the american heart association. *Circulation (New York, N.Y.)*. 2017;135(17):e927-e999. <https://www.ncbi.nlm.nih.gov/pubmed/28356445>. doi: 10.1161/CIR.0000000000000484.
2. Nakamura Y, Hama T, Nakamura Y, Tsukada H, Oda Y, Awa S. Orthostatic and exercise effects in children years after kawasaki disease. *Pediatr Cardiol*. 2020;41(1):24-30. <https://link.springer.com/article/10.1007/s00246-019-02216-x>. doi: 10.1007/s00246-019-02216-x.
3. Tuan S, Su H, Chen C, et al. Analysis of exercise capacity of children with kawasaki disease by a coronary artery z score model (ZSP version 4) derived by the lambda-mu-sigma method. *The Journal of pediatrics*. 2018;201:128-133. <https://dx.doi.org/10.1016/j.jpeds.2018.05.036>. doi: 10.1016/j.jpeds.2018.05.036.
4. Tuan S, Li M, Hsu M, et al. Cardiopulmonary function, exercise capacity, and echocardiography finding of pediatric patients with kawasaki disease. *Medicine*. 2016;95(2). https://explore.openaire.eu/search/publication?articleId=od_267::8345f4ac84d4143c83dd4b58235ee3bc. doi: 10.1097/MD.0000000000002444.
5. Chien K, Huang H, Weng K, et al. Arterial stiffness late after kawasaki disease in children: Assessment by performing brachial–ankle pulse wave velocity. *Journal of the Chinese Medical Association*. 2020;83(10):931. doi: 10.1097/jcma.0000000000000406.
6. Aggarwal V, Sexson-tejtal K, Lam W, et al. The incidence of arrhythmias during exercise stress tests among children with kawasaki disease: A single-center case series. *Congenital Heart Disease*. 2019;14(6):1032. doi: 10.1111/chd.12864.
7. Kuan T;, Chang Y;, Lin K, Chen G, Liou I, Tuan S. Differences between sexes in cardiopulmonary fitness among children and adolescents with kawasaki disease. *Healthcare*. 2022;10(2). doi: 10.3390/healthcare10020353.
8. Gravel H, Curnier D, Dallaire F, Fournier A, Portman M, Dahdah N. Cardiovascular response to exercise testing in children and adolescents late after kawasaki disease according to coronary condition upon onset. *Pediatr Cardiol*. 2015;36(7):1458-1464. <https://link.springer.com/article/10.1007/s00246-015-1186-5>. doi: 10.1007/s00246-015-1186-5.
9. Lin K, Liou I, Chen G, et al. Serial exercise testing and echocardiography findings of patients with kawasaki disease. *Front Pediatr*. 2022;10. doi: 10.3389/fped.2022.847343.
10. Yang T, Lee Y, Wang L, Chang T, Chang L, Kuo H. Patients with kawasaki disease have significantly low aerobic metabolism capacity and peak exercise load capacity during adolescence. *IJERPH*. 2020;17(22). doi: 10.3390/ijerph17228352.
11. Baker AL, Gauvreau K, Newburger JW, Sundel RP, Fulton DR, Jenkins KJ. Physical and psychosocial health in children who have had kawasaki disease. *Pediatrics (Evanston)*. 2003;111(3):579-583. <https://dx.doi.org/10.1542/peds.111.3.579>. doi: 10.1542/peds.111.3.579.
12. Banks L, Lin YT, Chahal N, Manlhiot C, Yeung RSM, McCrindle BW. Factors associated with low moderate-to-vigorous physical activity levels in pediatric patients with kawasaki disease. *Clinical pediatrics*. 2012;51(9):828-834.

<https://journals.sagepub.com/doi/full/10.1177/0009922812441664>. doi:
10.1177/0009922812441664.

13. Hansen K, Grady S, McCrindle BW, et al. Physicians' self-reported exercise testing and physical activity recommendations in kawasaki patients. *Pediatric cardiology*. 2022;1. <https://search.proquest.com/docview/2702192650>. doi: 10.1007/s00246-022-02984-z.
14. Gauthier N, Curran T, O'Neill JA, Alexander ME, Rhodes J. Establishing a comprehensive pediatric cardiac fitness and rehabilitation program for congenital heart disease. *Pediatr Cardiol*. 2020;41(8):1569-1579. <https://link.springer.com/article/10.1007/s00246-020-02413-z>. doi: 10.1007/s00246-020-02413-z.